arXiv:1608.07030v2 [math.CA] 30 Aug 2016

L_p-BOUNDS FOR THE ČEBYŠEV FUNCTIONAL

MOHAMMAD W. ALOMARI

Abstract. In this paper several new bounds for the Čebyšev functional involving L_p-norm are presented.

1. Introduction

In order to study the difference between two Riemann integral means, Barnett et al. [1] have proved the following estimates:

Theorem 1. *Let $f : [a,b] \to \mathbb{R}$ be an absolutely continuous function with the property that $f' \in L_\infty[a,b]$, i.e.,*

$$\|f'\|_\infty := ess \sup_{t\in[a,b]} |f'(t)|.$$

Then for $a \le c < d \le b$, we have the inequality

$$\begin{aligned}(1.1)\qquad &\left|\frac{1}{b-a}\int_a^b f(t)\,dt - \frac{1}{d-c}\int_c^d f(s)\,ds\right| \\ &\le \left[\frac{1}{4} + \left(\frac{(a+b)/2-(c+d)/2}{(b-a)-(d-c)}\right)^2\right][(b-a)-(d-c)]\,\|f'\|_\infty \\ &\le \frac{1}{2}[(b-a)-(d-c)]\,\|f'\|_\infty.\end{aligned}$$

The constant $1/4$ in the first inequality and $1/2$ in the second inequality are the best possible.

After that, Cerone and Dragomir [2] have obtained the following three results as well:

Theorem 2. *Let $f : [a,b] \to \mathbb{R}$ be an absolutely continuous mapping. Then for $a \le c < d \le b$, we have the inequalities*

$$\begin{aligned}(1.2)\qquad &\left|\frac{1}{b-a}\int_a^b f(t)\,dt - \frac{1}{d-c}\int_c^d f(s)\,ds\right| \\ &\le \begin{cases} \frac{(b-a)}{(q+1)^{1/q}}\left[1+\left(\frac{\rho}{1-\rho}\right)^q\right]^{1/q}\left[v^{q+1}+\lambda^{q+1}\right]^{1/q}\|f'\|_p, \\ \qquad\qquad f' \in L_p[a,b],\ 1 \le p < \infty,\ \frac{1}{p}+\frac{1}{q}=1; \\ \frac{1}{2}[1-\rho+|v-\lambda|]\,\|f'\|_1, \qquad f' \in L_1[a,b]; \end{cases}\end{aligned}$$

Date: August 31, 2016.
2010 *Mathematics Subject Classification.* 26D10, 26D15.
Key words and phrases. Čebyšev functional, absolutely continuous, norms.

where $(b-a)\,v = c-a$, $(b-a)\,\rho = d-c$ *and* $(b-a)\,\lambda = b-d$. *Both inequalities in (1.2) are sharp.*

In this paper several new error bounds for the Čebyšev functional involving L_p-norm are presented.

2. On Pečarić and Perić Result.

For the celebrated Čebyšev functional

$$\mathcal{T}(f,g) = \frac{1}{b-a}\int_a^b f(t)\,g(t)\,dt - \frac{1}{b-a}\int_a^b f(t)\,dt \cdot \frac{1}{b-a}\int_a^b g(t)\,dt. \tag{2.1}$$

the following sharp bounds involving L_p-norms are proved in literature:

Theorem 3. *Let* $f, g : [a,b] \to \mathbb{R}$ *be two absolutely continuous functions, then*

(2.2)

$$|\mathcal{T}(f,g)| \leq \begin{cases} \frac{(b-a)^2}{12}\|f'\|_\infty \|g'\|_\infty, & \text{if } f', g' \in L_\infty[a,b] \quad \text{proved in [3]} \\ \frac{(b-a)}{\pi^2}\|f'\|_2 \|g'\|_2, & \text{if } f', g' \in L_2[a,b] \quad \text{proved in [4]} \\ \frac{1}{8}(b-a)(M-m)\|g'\|_\infty, & \text{if } m \leq f \leq M,\ g' \in L_\infty[a,b] \ \text{ proved in [6]} \end{cases}$$

The constants $\frac{1}{12}$, $\frac{1}{\pi^2}$ *and* $\frac{1}{8}$ *are the best possible.*

In 2006, Pečarić and Perić [7] generalized the Čebyšev functional for n-times differentiable functions whose n-th derivatives are integrable. In special case, for $n = 1$ they obtained that:

Theorem 4. *(Pečarić & Perić (2006)) Let* $f, g : [0,1] \to \mathbb{R}$ *be two functions such that* $f' \in L_q[a,b]$ *and* $g' \in L_p[a,b]$ *are integrable on* $[0,1]$. *Then, the following inequalities are hold:*

$$|\mathcal{T}(f,g)| \leq \frac{1}{8}\|g'\|_p \|f'\|_q, \tag{2.3}$$

for all $p, q > 1$ *with* $\frac{1}{p} + \frac{1}{q} = 1$, *where* $\mathcal{T}(f,g)$ *is defined along the interval* $[0,1]$ *instead of* $[a,b]$.

and

$$|\mathcal{T}(f,g)| \leq \frac{1}{(q+1)^{\frac{1}{q}}}\left[\frac{\Gamma^2(p_1+1)}{\Gamma(2p_1+2)}\right]^{\frac{1}{p_1}}\|g'\|_p \|f'\|_{q_1}, \tag{2.4}$$

for all $p, q, p_1, q_1 > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$ and $\frac{1}{p_1} + \frac{1}{q_1} = 1$. Moreover, if one chooses

- $q_1 = p$, then (2.4) becomes

$$|\mathcal{T}(f,g)| \leq \frac{1}{(q+1)^{\frac{1}{q}}}\left[\frac{\Gamma^2(q+1)}{\Gamma(2q+2)}\right]^{\frac{1}{q}}\|g'\|_p \|f'\|_p. \tag{2.5}$$

- $q_1 = q$, then (2.4) becomes

$$|\mathcal{T}(f,g)| \leq \frac{1}{(q+1)^{\frac{1}{q}}}\left[\frac{\Gamma^2(p+1)}{\Gamma(2p+2)}\right]^{\frac{1}{p}}\|g'\|_p \|f'\|_q. \tag{2.6}$$

In particular, they presented an optimal value of $C_{a,b}(p,q)$ for arbitrary conjugate exponents p,q, i.e., $\frac{1}{p}+\frac{1}{q}=1$, $p>1$. Namely, if $f'\in\in L_p[a,b]$ and $g'\in L_q[a,b]$, then

$$|\mathcal{T}(f,g)|\leq\frac{1}{8}\left\|f'\right\|_p\left\|g'\right\|_q,\quad\forall\ p,q>1,\ \frac{1}{p}+\frac{1}{q}=1. \tag{2.7}$$

Unfortunately, wrong calculations had been made in [7]. Let us observe that the function presented in [7], does not satisfies the optimal sharpness where the authors seeking the best possibility of $\frac{1}{8}$ in the case $p=\infty,q=1$.

Example 1. *Let $g(x)=x$, $x\in[0,1]$ and $\widetilde{f}_\epsilon(x):=\int_0^x\widetilde{f}_\epsilon(t)dt$, where*

$$\widetilde{f}_\epsilon(t)=\begin{cases}\frac{1}{2\cdot\epsilon}, & \epsilon\in\left(\frac{1}{2}-\epsilon,\frac{1}{2}+\epsilon\right)\\ 0, & \text{otherwise}\end{cases}$$

for all $\epsilon\in\left(0,\frac{1}{2}\right)$. It is not difficult to compute

$$\begin{aligned}\widetilde{f}_\epsilon(x):=\int_0^x\widetilde{f}_\epsilon(t)dt&=\int_0^{\frac{1}{2}-\epsilon}\widetilde{f}_\epsilon(t)dt+\int_{\frac{1}{2}-\epsilon}^x\widetilde{f}_\epsilon(t)dt\\&=0+\int_{\frac{1}{2}-\epsilon}^x\frac{1}{2\epsilon}dt\\&=\left(x-\frac{1}{2}+\epsilon\right)\frac{1}{2\epsilon},\qquad x\in[0,1]\end{aligned}$$

$\int_0^1\widetilde{f}_\epsilon(x)dx=\frac{1}{2\epsilon}\int_0^1\left(x-\frac{1}{2}+\epsilon\right)dx=\frac{\left(\frac{1}{2}+\varepsilon\right)^2-\left(-\frac{1}{2}+\varepsilon\right)^2}{4\varepsilon}$, $\int_0^1 xdx=\frac{1}{2}$, $\int_0^1\widetilde{f}_\epsilon(x)g(x)dx=\frac{1}{2\epsilon}\int_0^1 x\left(x-\frac{1}{2}+\epsilon\right)dx=\frac{1}{24\epsilon}+\frac{1}{4}$

$$\left|\frac{1}{24\epsilon}+\frac{1}{4}-\frac{\left(\frac{1}{2}+\varepsilon\right)^2-\left(-\frac{1}{2}+\varepsilon\right)^2}{8\varepsilon}\right|\leq C\cdot\frac{1}{2\epsilon}\cdot 1$$

$$\lim_{\varepsilon\to0^+}\left|\frac{1}{12}+\frac{\epsilon}{2}-\frac{\left(\frac{1}{2}+\epsilon\right)^2-\left(-\frac{1}{2}+\epsilon\right)^2}{4}\right|=\frac{1}{12}\leq C.$$

Hence, the suggested functions does not satisfy the sharpness of (2.7) in the case $p=\infty,q=1$, as claimed in [7].

We remark that the first result involving absolutely continuous functions whose first derivatives belong to L_p spaces was unpublished and not achieved as cited by the authors of [5]. This result was presented by Beesack, Mitrinović and Vasić and mentioned in [5], (p.302) without proof, which reads:

$$|\mathcal{T}(f,g)|\leq\frac{b-a}{4}\left(\frac{2^p-1}{p(p+1)}\right)^{\frac{1}{p}}\left(\frac{2^q-1}{q(q+1)}\right)^{\frac{1}{q}}\left\|f'\right\|_p\left\|g'\right\|_q, \tag{2.8}$$

where $\|h\|_p:=\left(\int_a^b|h(t)|^p\,dt\right)^{1/p}$, $\forall p\geq1$ and $\frac{1}{p}+\frac{1}{q}=1$.

We remark here, for the constant

$$\omega(p,q):=\frac{1}{4}\left(\frac{2^p-1}{p(p+1)}\right)^{1/p}\left(\frac{2^q-1}{q(q+1)}\right)^{1/q},$$

we have

$$\frac{1}{8} \leq \omega(p,q) \leq \frac{1}{4},$$

for all $q = \frac{p}{p-1}$, $p > 1$. Furthermore, we have the following particular cases in (2.8).

(1) If $p = q = 2$, we have

$$|\mathcal{T}(f,g)| \leq \frac{b-a}{8} \|f'\|_2 \|g'\|_2 . \tag{2.9}$$

(2) If $q \longrightarrow \infty$, we have

$$|\mathcal{T}(f,g)| \leq \frac{b-a}{4} \|f'\|_1 \|g'\|_\infty . \tag{2.10}$$

The first case was improved by Lupaş [4] and the second was improved in [7] (and in this work as well). As a final comment, the problem that what is the best possible constant $C_{a,b}(p,q)$ would satisfies the inequality

$$|\mathcal{T}(f,g)| \leq C_{a,b}(p,q) \|f'\|_p \|g'\|_q , \tag{2.11}$$

for all $p,q > 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$? is an open problem.

3. The Results

Another proof for the Čebyšev inequality (the first inequality in(2.2)) is considered as follows:

Theorem 5. *Let $f, g : [a,b] \to \mathbb{R}$ be two absolutely continuous on $[a,b]$. If f' is bounded on $[a,b]$, i.e., $\|f'\|_\infty := \sup_{t\in[a,b]} |f'(t)| < \infty$, then*

$$|\mathcal{T}(f,g)| \leq \frac{(b-a)^2}{12} \|g'\|_\infty \|f'\|_\infty \tag{3.1}$$

The constant $\frac{1}{12}$ is the best possible.

Proof. Using integration by parts, we have

$$\mathcal{T}(f,g) = -\frac{1}{b-a}\int_a^b \left(\int_a^t g(s)\,ds - \frac{t-a}{b-a}\int_a^b g(s)\,ds\right) f'(t)\,dt. \tag{3.2}$$

Taking the absolute value in (3.2) and utilizing the triangle inequality, so as f' is in $L_\infty([a,b])$, we have

$$\begin{aligned}
|\mathcal{T}(f,g)| &\leq \frac{1}{b-a}\int_a^b \left|(t-a)\left[\frac{1}{t-a}\int_a^t g(u)\,du - \frac{1}{b-a}\int_a^b g(u)\,du\right]\right| |f'(t)|\,dt \\
&\leq \frac{\|f'\|_\infty}{b-a}\cdot \int_a^b (t-a)\left|\frac{1}{t-a}\int_a^t g(u)\,du - \frac{1}{b-a}\int_a^b g(u)\,du\right| dt \qquad (3.3)\\
&\leq \frac{\|f'\|_\infty}{2(b-a)} \|g'\|_\infty \int_a^b (t-a)(b-t)\,dt \\
&= \frac{1}{12}(b-a)^2 \|f'\|_\infty \|g'\|_\infty ,
\end{aligned}$$

where, in (3.3) we have used the inequality (1.1), with $d = t$ and $c = a$. The sharpness holds with the identity function $f(x) = x$, $x \in [a,b]$. □

Theorem 6. *Let $f, g : [a, b] \to \mathbb{R}$ be two absolutely continuous on $[a, b]$. If f' is bounded on $[a, b]$, i.e., $\|f'\|_\infty := \sup_{t\in[a,b]} |f'(t)| < \infty$ and $g' \in L_p[a, b]$ then*

$$|\mathcal{T}(f,g)| \le \frac{(b-a)^2}{(q+1)^{1/q}} \mathrm{B}\left(2, 1+\frac{1}{q}\right) \cdot \|g'\|_p \|f'\|_\infty \tag{3.4}$$

where, $p > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. To prove (3.4), in (3.3) we apply (1.2), by setting $d = t$ and $c = a$, we get

$$\left|\frac{1}{t-a}\int_a^t g(s)\,ds - \frac{1}{b-a}\int_a^b g(s)\,ds\right| \le \frac{(b-t)^{\frac{1}{q}}}{(q+1)^{1/q}(b-a)^{\frac{1}{q}}}\left[(t-a)^q + (b-t)^q\right]^{1/q}\|g'\|_p \tag{3.5}$$

for all $1 < p < \infty$, $\frac{1}{p} + \frac{1}{q} = 1$.

Substituting (3.5) in (3.3), we get

$$\begin{aligned}
|\mathcal{T}(f,g)| &\le \frac{\|f'\|_\infty}{(b-a)} \frac{\|g'\|_p}{(q+1)^{1/q}(b-a)^{1/q}} \int_a^b (t-a)(b-t)^{\frac{1}{q}}\left[(t-a)^q + (b-t)^q\right]^{1/q} dt \\
&\le \frac{\|f'\|_\infty}{(b-a)} \frac{\|g'\|_p}{(q+1)^{1/q}(b-a)^{1/q}} \sup_{t\in[a,b]}\left[(t-a)^q + (b-t)^q\right]^{1/q} \int_a^b (t-a)(b-t)^{\frac{1}{q}}\,dt \\
&= \|f'\|_\infty \frac{1}{(q+1)^{1/q}}(b-a)^2\,\mathrm{B}\left(2, 1+\frac{1}{q}\right)\cdot\|g'\|_p
\end{aligned}$$

which proves (3.4). □

For two absolutely continuous functions whose first derivatives belongs to $L_1[a, b]$, a sharp inequality is incorporated in the following theorem.

Theorem 7. *Let $f, g : [a, b] \to \mathbb{R}$ be two absolutely continuous on $[a, b]$. If $f' \in L_1[a, b]$, then*

$$|\mathcal{T}(f,g)| \le \|f'\|_1 \begin{cases} \frac{(b-a)}{4(q+1)^{1/q}} \cdot \|g'\|_p, & \text{if } g' \in L_p[a,b]; \\ \frac{1}{4}\|g'\|_1, & \text{if } g' \in L_1[a,b]; \end{cases} \tag{3.6}$$

where, $p > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. As $f' \in L_1([a, b])$, we have

$$\begin{aligned}
|\mathcal{T}(f,g)| &\le \frac{1}{b-a}\int_a^b \left|(t-a)\left[\frac{1}{t-a}\int_a^t g(u)\,du - \frac{1}{b-a}\int_a^b g(u)\,du\right]\right| |f'(t)|\,dt \\
&\le \frac{1}{b-a}\sup_{t\in[a,b]}(t-a)\left|\frac{1}{t-a}\int_a^t g(u)\,du - \frac{1}{b-a}\int_a^b g(u)\,du\right| \cdot \int_a^b |f'(t)|\,dt. \qquad (3.7)
\end{aligned}$$

But

$$\frac{1}{b-a}\sup_{t\in[a,b]}(t-a)\left|\frac{1}{t-a}\int_a^t g(u)\,du-\frac{1}{b-a}\int_a^b g(u)\,du\right|$$

$$\leq\frac{1}{(b-a)}\begin{cases}\frac{\|g'\|_p}{(q+1)^{1/q}(b-a)^{1/q}}\sup\limits_{t\in[a,b]}\left\{(t-a)(b-t)^{\frac{1}{q}}\left[(t-a)^q+(b-t)^q\right]^{1/q}\right\}, & g'\in L_p[a,b],\\ \frac{\|g'\|_1}{b-a}\sup\limits_{t\in[a,b]}(t-a)(b-t), & g'\in L_1[a,b],\end{cases}$$

$$=\begin{cases}\frac{(b-a)}{4(q+1)^{1/q}}\cdot\|g'\|_p, & g'\in L_p[a,b],\\ \frac{1}{4}\|g'\|_1, & g'\in L_1[a,b],\end{cases}$$

which proves the first and the second inequalities in (3.6). □

Theorem 8. *Let $f,g:[a,b]\to\mathbb{R}$ be two absolutely continuous on $[a,b]$. If $f'\in L_\alpha[a,b]$, $\alpha>1$, $\frac{1}{\alpha}+\frac{1}{\beta}=1$, then*

(3.8) $|\mathcal{T}(f,g)|$

$$\leq\begin{cases}\frac{(b-a)^{\frac{1}{\beta}}}{2}\cdot \mathrm{B}^{\frac{1}{\beta}}(\beta+1,\beta+1)\cdot\|f'\|_\alpha\cdot\|g'\|_\infty, & \text{if } g'\in L_\infty[a,b]\\ \frac{(b-a)^{1+\frac{1}{\beta}}}{(q+1)^{1/q}}\cdot \mathrm{B}^{\frac{1}{\beta}}\left(\beta+1,\frac{\beta}{q}+1\right)\|g'\|_p\|f'\|_\alpha\cdot\|g'\|_p, & \text{if } g'\in L_p[a,b],\\ & p>1,\ \frac{1}{p}+\frac{1}{q}=1;\\ (b-a)^{\frac{1}{\beta}}\cdot \mathrm{B}^{\frac{1}{\beta}}(\beta+1,\beta+1)\cdot\|f'\|_\alpha\cdot\|g'\|_1, & \text{if } g'\in L_1[a,b]\end{cases}$$

Proof. Taking the absolute value in (3.2) and utilizing the triangle inequality. As $f'\in L_\alpha([a,b])$, we have

(3.9)

$$|\mathcal{T}(f,g)|\leq\frac{1}{b-a}\int_a^b\left|(t-a)\left[\frac{1}{t-a}\int_a^t g(u)\,du-\frac{1}{b-a}\int_a^b g(u)\,du\right]\right||f'(t)|\,dt$$

$$\leq\frac{1}{b-a}\left(\int_a^b|t-a|^\beta\left|\frac{1}{t-a}\int_a^t g(u)\,du-\frac{1}{b-a}\int_a^b g(u)\,du\right|^\beta dt\right)^{1/\beta}\left(\int_a^b|f'(t)|^\alpha\,dt\right)^{1/\alpha}$$

Now, in (1.1) put $d=t$ and $c=a$, then

$$\left|\frac{1}{t-a}\int_a^t g(u)\,du-\frac{1}{b-a}\int_a^b g(u)\,du\right|\leq\frac{b-t}{2(b-a)}\cdot\|g'\|_\infty$$

Substituting in (3.10) we get

$$\begin{aligned}|\mathcal{T}(f,g)|&\leq\frac{1}{2(b-a)^2}\cdot\|f'\|_\alpha\cdot\|g'\|_\infty\left(\int_a^b(t-a)^\beta(b-t)^\beta\,dt\right)^{\frac{1}{\beta}}\\&=\frac{(b-a)^{\frac{1}{\beta}}}{2}\cdot\mathrm{B}^{\frac{1}{\beta}}(\beta+1,\beta+1)\cdot\|f'\|_\alpha\cdot\|g'\|_\infty\end{aligned}$$

which prove the first inequality in (3.8).

To prove the second and third inequalities in (3.8), we apply (1.2), by setting $d=t$ and $c=a$, we get

$$\int_a^b|t-a|^\beta\left|\frac{1}{t-a}\int_a^t g(u)\,du-\frac{1}{b-a}\int_a^b g(u)\,du\right|^\beta dt \tag{3.10}$$

$$\leq\begin{cases}\dfrac{\|g'\|_p^\beta}{(q+1)^{\beta/q}(b-a)^{\frac{\beta}{q}}}\int_a^b(t-a)^\beta(b-t)^{\frac{\beta}{q}}\left[(t-a)^q+(b-t)^q\right]^{\beta/q}dt,\\ \qquad\qquad\text{if } g'\in L_p[a,b],\ \beta>1,\ 1<p<\infty,\ \frac{1}{p}+\frac{1}{q}=1;\\ \dfrac{1}{(b-a)^\beta}\|g'\|_1^\beta\cdot\int_a^b(t-a)^\beta(b-t)^\beta,\quad\text{if } g'\in L_1[a,b].\end{cases}$$

$$\leq\begin{cases}\dfrac{\|g'\|_p^\beta}{(q+1)^{\beta/q}(b-a)^{\frac{\beta}{q}}}\sup\limits_{t\in[a,b]}\left[(t-a)^q+(b-t)^q\right]^{\beta/q}\int_a^b(t-a)^\beta(b-t)^{\frac{\beta}{q}}dt,\\ \qquad\qquad\text{if } g'\in L_p[a,b],\ \beta>1,\ 1<p<\infty,\ \frac{1}{p}+\frac{1}{q}=1;\\ \dfrac{1}{(b-a)^\beta}\|g'\|_1^\beta\cdot(b-a)^{2\beta+1}\mathrm{B}(\beta+1,\beta+1),\quad\text{if } g'\in L_1[a,b].\end{cases}$$

$$=\begin{cases}\dfrac{(b-a)^{\left(2+\frac{1}{q}\right)\beta+1}}{(q+1)^{\beta/q}(b-a)^{\frac{\beta}{q}}}\cdot\mathrm{B}\left(\beta+1,\frac{\beta}{q}+1\right)\|g'\|_p^\beta,\\ \qquad\qquad\text{if } g'\in L_p[a,b],\ \beta>1,\ p>1,\ \frac{1}{p}+\frac{1}{q}=1;\\ \dfrac{1}{(b-a)^\beta}\|g'\|_1^\beta\cdot(b-a)^{2\beta+1}\mathrm{B}(\beta+1,\beta+1),\quad\text{if } g'\in L_1[a,b].\end{cases}$$

Substituting (3.10) in (3.9), we get

$$|\mathcal{T}(f,g)|$$

$$\leq\frac{\|f'\|_\alpha}{(b-a)}\begin{cases}\dfrac{(b-a)^{2+\frac{1}{\beta}}}{(q+1)^{1/q}}\cdot\mathrm{B}^{\frac{1}{\beta}}\left(\beta+1,\frac{\beta}{q}+1\right)\|g'\|_p,\\ \qquad\qquad\text{if } g'\in L_p[a,b],\ \beta>1,\ p>1,\ \frac{1}{p}+\frac{1}{q}=1;\\ (b-a)^{1+\frac{1}{\beta}}\cdot\mathrm{B}^{\frac{1}{\beta}}(\beta+1,\beta+1)\cdot\|g'\|_1,\quad\text{if } g'\in L_1[a,b].\end{cases}$$

which proves the second and the third inequalities in (3.8). □

Remark 1. *In (3.8) we have the following particular cases*

(1) *If $\alpha = p$ and $\beta = q$, then we have*

$$|\mathcal{T}(f,g)| \leq \frac{(b-a)^{1+\frac{1}{q}}}{(q+1)^{1/q}} \cdot \mathrm{B}^{\frac{1}{q}}(q+1,2) \left\|f'\right\|_p \cdot \left\|g'\right\|_p .$$

(2) *If $\alpha = q$ and $\beta = p$, then we have*

$$|\mathcal{T}(f,g)| \leq \frac{(b-a)^{1+\frac{1}{p}}}{(q+1)^{1/q}} \cdot \mathrm{B}^{\frac{1}{p}}\left(p+1, \frac{p}{q}+1\right) \left\|f'\right\|_q \cdot \left\|g'\right\|_p . \tag{3.11}$$

In (3.11), as $q = 1$ and $p = \infty$,

$$|\mathcal{T}(f,g)| \leq \frac{b-a}{8} \left\|f'\right\|_1 \left\|g'\right\|_\infty .$$

which improves (2.10).

References

[1] N.S. Barnett, P. Cerone, S.S. Dragomir and A.M. Fink, Comparing two integral means for absolutely continuous mappings whose derivatives are in $L_\infty[a,b]$ and applications, *Comp. and Math. Appl.*, 44 (1/2) (2002), 241–251.

[2] P. Cerone and S.S. Dragomir, Differences between means with bounds from a Riemann–Stieltjes integral, *Comp. Math. Appl.*, 46 (2003) 445–453.

[3] P.L. Čebyšev, Sur les expressions approximatives des intègrals dèfinis par les outres prises entre les même limites, *Proc. Math. Soc. Charkov*, 2 (1882), 93–98.

[4] A. Lupaş, The best constant in an integral inequality, *Mathematica*, (Cluj, Romania) 15 (38)(2) (1973), 219–222.

[5] D.S. Mitrinović, J.E. Pečarić and A.M. Fink, Classical and New Inequalities in Analysis, Kluwer Academic Publishers, Dordrecht, 1993.

[6] A.M. Ostrowski, On an integral inequality, *Aequat. Math.*, 4 (1970), 358–373.

[7] J. Pečarić and I. Perić, Identities for the Chebyshev functional involving derivatives of arbitrary order and applications, *J. Math. Anal. Appl.*, 313 (2006), 475–483.

Department of Mathematics, Faculty of Science and Information Technology, Irbid National University, 21110 Irbid, Jordan

E-mail address: mwomath@gmail.com